



Review Article

Aging with HIV: Triumphs and Challenges

Alice Zhabokritsky^{1,2} and Sharon L Walmsley^{1,2,3*}

¹Department of Medicine, University Health Network, Toronto, Canada

²Department of Medicine, University of Toronto, Toronto, Canada

³Toronto General Hospital Research Institute, University Health Network, Toronto, Canada

Abstract

The HIV population is aging. Antiretroviral therapy has significantly improved the life expectancy of people living with HIV, which is now approaching that of the general population. However, older adults with HIV are at a greater risk of developing comorbidities and experiencing polypharmacy, and they tend to encounter these challenges at a younger age. There is an ongoing need to adapt healthcare infrastructure and public health policies to address the complex needs of the aging HIV population.

Keywords: Aging; Comorbidity; HIV; Polypharmacy; Prevention; Women

Abbreviations

HIV: Human Immunodeficiency Virus
AIDS: Acquired Immunodeficiency Syndrome
PLWH: People Living With HIV
LTS: Long Term Survivors
CYPP450: Cytochrome P450
LTC: Long-term Care

Introduction

HIV care has gone through an amazing transformation over the past 30 years. What was once a terminal diagnosis can now be managed as a chronic condition. People are living with HIV longer than ever before as treatment regimens become more effective, tolerable and initiated earlier in the disease progress at higher CD4 counts.

***Corresponding author:** Sharon L Walmsley, Department of Medicine, University Health Network, 200 Elizabeth Street, Room 13EN214, Toronto, M5G2C4, Canada, Tel: +1 4163403871; E-mail: sharon.walmsley@uhn.ca

Citation: Zhabokritsky A, Walmsley SL (2020) Aging with HIV: Triumphs and Challenges. J Gerontol Geriatr Med 6: 062.

Received: July 17, 2020; **Accepted:** July 27, 2020; **Published:** August 03, 2020

Copyright: © 2020 Zhabokritsky A, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Today nearly half of People Living With HIV (PLWH) in North America are over the age 50 [1,2], with rising numbers of older adults living with HIV around the world [3]. As we celebrate this changing demographic, it is important to acknowledge the challenges faced by this aging population and adapt our clinical and public health practices accordingly.

Diversity Among Older Adults with HIV

Firstly, it is important to recognize that older adults with HIV represent a diverse population of individuals that may face unique challenges related to their physical and mental health. For example, Long Term Survivors (LTS), individuals who were diagnosed with HIV before effective antiretroviral therapy became available [4], received some of the earliest HIV therapies, exposing them to significant toxicities and increasing their risk of developing chronic conditions such as lipodystrophy, diabetes, peripheral neuropathy and metabolic syndrome. Many LTS have experienced immense psychosocial trauma related to receiving a diagnosis that was once considered a death sentence, losing friends and loved ones to the condition, facing years of stigma, poverty, food insecurity and social isolation [5]. Providing trauma-informed care in a multidisciplinary setting may be of particular benefit for this group.

In contrast, there are also a substantial proportion of older adults with a recent HIV diagnosis. In 2018, people over the age of 50 accounted for 22.5% of new HIV diagnoses in Canada [1] and similarly in the United States, 1 in 6 new diagnoses in 2017 were among those over age 50 [2]. These numbers include individuals that recently acquired HIV as well as those that were recently tested but acquired the infection earlier. Individuals with delayed diagnosis have more advanced immune dysregulation at the time of presentation and are at greater risk of presenting with opportunistic infections and comorbidities related to chronic uncontrolled HIV infection. On the other hand, older adults with recently acquired HIV infection may have relatively preserved immune systems but require more guidance in terms of access to HIV resources and community supports. Finally, the mode of HIV transmission might carry additional information about the social aspects of an individual's diagnosis and journey, such as experiences related to sexual identity, drug use or sex work. Undoubtedly, it is important to recognize the multifaceted needs of older adults with HIV and have an appreciation for their varied backgrounds.

Delayed HIV Diagnosis in Older Adults and Women

Older adults are more likely to have unrecognized HIV infection for many years and subsequently be diagnosed as inpatients, at lower CD4 counts and at more advanced stages of HIV infection relative to younger adults [6,7]. Women are particularly at increased risk of being diagnosed late in their illness and are less likely to be tested in the outpatient setting [8]. Delayed diagnosis in women and older adults reflects low perceived risk of HIV infection among patients and their healthcare providers. Late treatment initiation is especially detrimental in the aging population. In 2015, a large international

randomised controlled trial showed a clear morbidity and mortality benefit to early treatment initiation at high baseline CD4 counts, compared to delaying treatment until CD4 count drops below 350, across age groups [9]. However, rates of AIDS-defining illnesses, serious non-AIDS illnesses (including cardiovascular disease, end-stage renal and liver disease and non-AIDS cancers) and death were disproportionately higher in older adults if treatment was delayed. Absolute risk reduction was greatest in the >50 age group with the lowest number needed to treat to prevent an adverse clinical event, demonstrating that older adults benefit the most from early recognition and treatment of HIV infection. These findings contributed to a change in practice, which now emphasizes early treatment initiation, at high CD4 counts. Unrecognized infection in older adults prevents them from benefiting from this treatment approach. Improving risk factor recognition along with implementation of routine HIV screening are therefore of great importance to improve survival and reduce morbidity in this population.

Increasing Life Expectancy but not Comorbidity-Free Years

The life expectancy of PLWH continues to increase and is approaching that of the general population. In North America, total life expectancy at age 20 increased from 56.1 years in 2000-2002 to 71.4 years in 2006-2007 among PLWH [10]. In contrast, life expectancy in the general population in Canada in 2007 was 80.5, and 78 in the United States [11,12]. The life expectancy gap relative to the general population narrowed further to just 3.4 years in 2014-2016 in an American cohort of PLWH with early treatment initiation [13]. Although people are living longer with HIV, they tend to spend a substantially greater number of years dealing with chronic conditions with fewer healthy years free of comorbidities. On average, PLWH developed common medical comorbidities 16 years earlier than those in the general population [13]. This number has not improved since early 2000's or with implementation of early treatment strategies.

Comorbidities among PLWH

PLWH are at a substantially increased risk relative to the general population of developing medical comorbidities, including cardiovascular disease, diabetes, renal failure and liver disease, and they are more likely to experience polypharmacy and drug-drug interactions [14-16]. The difference in risk increases as people age. Co-infection with cytomegalovirus seems to be an independent risk factor for developing non-AIDS defining comorbidities, especially cardiovascular and cerebrovascular diseases [17]. Prevalence of osteopenia and osteoporosis is also substantially higher among PLWH with greater risk of bone fracture across age groups [14,18]. Certain cancers are more prevalent among PLWH, including AIDS-defining (i.e., Kaposi's sarcoma, non-Hodgkin's lymphoma, cervical cancer) and non-AIDS defining cancers (i.e., Hodgkin's lymphoma, lung, liver, anal and oropharyngeal cancers) [19,20]. With advancing age and time since HIV diagnosis, cancer burden shifts from AIDS-defining to non-AIDS defining cancers [19]. This change is a reflection of HIV treatment initiation and recovery of the CD4 count, coupled with prolonged exposure to risk factors, such as smoking, alcohol consumption and co-infections with human papillomavirus and Hepatitis C virus. Smoking is highly prevalent among PLWH (>50% report active smoking), with rates that are three times that of the general population [21,22]. People with HIV are also less likely

to quit smoking [21], contributing substantially to development of comorbid disease.

In order to reduce the burden related to co-morbidities commonly experienced by PLWH, routine screening and utilization of preventative strategies is required. These should include assessments of blood pressure, glycemic control, lipids profile, renal function and cancer screening, along with testing for co-infections and vaccination. Smoking cessation programs are critical to address the large burden of tobacco dependence in this population. An approach that combines behavioural and pharmacological treatments to facilitate smoking cessation has been found to be more successful than standalone interventions and should be integrated into routine HIV care [22,23].

Finally, comorbidity management must implement careful medication review to avoid potential drug-drug interactions. As people age with HIV, they are more likely to be taking medications in addition to anti-retroviral therapy, such as cardiovascular medications, lipid-lowering agents, antiplatelets, anticoagulants, acid-lowering medications and hypoglycemics, as well as psychoactive medications and analgesics [24,25]. Many anti-retroviral agents are metabolized through the Cytochrome P450 (CYP450) enzyme system [24]. They can also exert bidirectional effects on CYP450 isoenzymes, potentially resulting in increased or decreased metabolism of many commonly prescribed medications. On the other hand, medications that impact absorption or metabolism of anti-retroviral agents can impact treatment efficacy. Given the complex nature of these interactions and increased potential for toxicities, clinicians need to routinely screen for drug-drug interactions with guidance from online resources [26] or a pharmacist with experience in HIV treatment.

Women's Health

Women aging with HIV may encounter some additional challenges. For example, they may experience menopause at an earlier age relative to the general population and experience a higher burden of symptoms and metabolic changes associated with menopause [27-29]. In particular, psychological and vasomotor symptoms have been reported to be more common in women living with HIV [29]. As women with HIV transition through menopause, they face higher rates of osteoporosis, cardiovascular and cerebral vascular disease and impaired lipid and glucose metabolism [30,31]. Furthermore, symptoms related to HIV infection or its treatment can be mistaken for menopause, such as night-sweats, sleep disturbances, menstrual cycle irregularities or vaginal dryness [27]. Conditions associated with HIV infection, including tuberculosis and lymphoma, can present with symptoms similar to menopause and should be considered in the right clinical context. Finally, it is important for clinicians to recognize that as women age, they may be at an elevated risk for HIV transmission due to thinning of the vaginal and cervical mucosa with menopause, immunological changes in the cervix, and reduced condom use when contraception is no longer a concern [32,33].

Geriatric Syndromes

Geriatric syndromes are multifactorial conditions used to identify vulnerable older adults and include conditions such as falls, frailty, functional and cognitive impairment [34]. Traditional risk factors for developing geriatric syndromes include psychosocial factors such as social isolation and substance use, as well as medical factors including multimorbidity, polypharmacy and chronic inflammation -

factors that are common in the aging HIV population. Consequently, geriatric syndromes among PLWH have been reported to occur at the same rates but several years earlier than in the general population [34]. Lower nadir CD4 count was associated with increased risk of developing geriatric syndromes, revealing a potential target for intervention unique to PLWH in addition to traditional risk factor modification. As the aging HIV population grows, we need to consider how to deliver comprehensive care to this population, including assessment, prevention and management of geriatric syndromes.

Long-Term Healthcare Facilities

PLWH currently represent a small proportion of individuals living in Long-Term Care (LTC) or complex continuing care facilities, however, as the HIV population ages these numbers are expected to increase [35]. In general, healthcare providers working in these facilities have little experience caring for PLWH. It has been recognized that managing comorbidities, mental health issues and polypharmacy among PLWH entering LTC facilities may be especially challenging and services will need to be adapted to address the needs of this population [36]. Furthermore, issues of stigma can be amplified as individuals enter LTC facilities. Many will experience anticipated and enacted stigma as they reveal their HIV status while trying to create new social networks, negatively affecting their psychological well-being and contributing to increased social isolation [37]. Institutional policies that empower PLWH to maintain their existing social supports, which are often informal and fragile, as they enter LTC, can be protective against the effects of stigma [36,38]. There are ongoing efforts to educate LTC providers about the unique needs of PLWH and the expected progression of illness, with the goal of promoting compassionate and informed care for this population [39].

Conclusion

People are living longer with HIV but they tend to develop medical comorbidities earlier and more frequently than those in the general population, resulting in fewer healthy years. Delayed treatment initiation is especially detrimental in the aging HIV population and therefore early detection needs to be prioritized through screening. As people age, preventative care and risk factor modification become integral components of HIV care in addition to management of geriatric syndromes.

Funding

No external funding.

Conflicts of Interest

The authors declare no conflicts of interest.

References

- Haddad N, Robert A, Weeks A, Popovic N, Siu W, et al. (2018) HIV in Canada-Surveillance Report, 2018. *Can Commun Dis Rep* 45: 304-312.
- Centers for Disease Control and Prevention (2018) Diagnoses of HIV Infection in the United States and Dependent Areas, 2018. HIV Surveillance Report, CDC, Georgia, USA.
- Autenrieth CS, Beck EJ, Stelzle D, Mallouris C, Mahy M, et al. (2018) Global and regional trends of people living with HIV aged 50 and over: Estimates and projections for 2000-2020. *PLoS One* 13: 0207005.
- Krentz HB, Gill MJ (2018) Long-term HIV/AIDS survivors: Patients living with HIV infection retained in care for over 20 years. What have we learned? *Int J STD AIDS* 29: 1098-1105.
- Broun SN (1998) Understanding "post-AIDS survivor syndrome": A record of personal experiences. *AIDS Patient Care STDS* 12: 481-488.
- Ellman TM, Sexton ME, Warshafsky D, Sobieszczyk ME, Morrison EA (2014) A forgotten population: Older adults with newly diagnosed HIV. *AIDS Patient Care STDS* 28: 530-536.
- Iwuji CC, Churchill D, Gilleece Y, Weiss HA, Fisher M (2013) Older HIV-infected individuals present late and have a higher mortality: Brighton, UK cohort study. *BMC Public Health* 13: 397.
- Mugavero MJ, Castellano C, Edelman D, Hicks C (2007) Late diagnosis of HIV infection: The role of age and sex. *Am J Med* 120: 370-373.
- Molina JM, Grund B, Gordin F, Williams I, Schechter M, et al. (2018) Which HIV-infected adults with high CD4 T-cell counts benefit most from immediate initiation of antiretroviral therapy? A post-hoc subgroup analysis of the START trial. *Lancet HIV* 5: 172-180.
- Samji H, Cescon A, Hogg RS, Modur SP, Althoff KN, et al. (2013) Closing the gap: Increases in life expectancy among treated HIV-positive individuals in the United States and Canada. *PLoS One* 8: 81355.
- Statistics Canada (2020) Life expectancy at birth and at age 65, by province and territory, three-year average. Statistics Canada, Ottawa, Canada.
- Xu JQ, Kochanek KD, Tejada-Vera B (2009) Deaths: preliminary data for 2007. *Natl Vital Stat Rep* 58.
- Marcus JL, Leyden W, Anderson AN, Hechter R, Horberg MA, et al. (2020) Increased overall life expectancy but not comorbidity-free years for people with HIV. CROI, San Francisco, USA.
- Guaraldi G, Orlando G, Zona S, Menozzi M, Carli F, et al. (2011) Premature age-related comorbidities among HIV-infected persons compared with the general population. *Clin Infect Dis* 53: 1120-1126.
- Feinstein MJ, Hsue PY, Benjamin LA, Bloomfield GS, Currier JS, et al. (2019) Characteristics, Prevention, and Management of Cardiovascular Disease in People Living With HIV: A Scientific Statement From the American Heart Association. *Circulation* 140: 98-124.
- Tseng A, Szadkowski L, Walmsley S, Salit I, Raboud J (2013) Association of age with polypharmacy and risk of drug interactions with antiretroviral medications in HIV-positive patients. *Ann Pharmacother* 47: 1429-1439.
- Lichtner M, Cicconi P, Vita S, Cozzi-Lepri A, Galli M, et al. (2015) Cytomegalovirus coinfection is associated with an increased risk of severe non-AIDS-defining events in a large cohort of HIV-infected patients. *J Infect* 211: 178-186.
- Bonjoch A, Figueras M, Estany C, Perez-Alvarez N, Rosales J, et al. (2010) High prevalence of and progression to low bone mineral density in HIV-infected patients: A longitudinal cohort study. *AIDS* 24: 2827-2833.
- Robbins HA, Pfeiffer RM, Shiels MS, Li J, Hall HI, et al. (2015) Excess cancers among HIV-infected people in the United States. *J Natl Cancer Inst* 107: dju503.
- Hernández-Ramírez RU, Shiels MS, Dubrow R, Engels EA (2017) Cancer risk in HIV-infected people in the USA from 1996 to 2012: A population-based, registry-linkage study. *Lancet HIV* 4: 495-504.
- Mdodo R, Frazier EL, Dube SR, Mattson CL, Sutton MY, et al. (2015) Cigarette smoking prevalence among adults with HIV compared with the general adult population in the United States: Cross-sectional surveys. *Ann Intern Med* 162: 335-344.
- Rapid Response Service (2015) Smoking Cessation Interventions in People with HIV/AIDS. Rapid Response Service.

23. Ledgerwood DM, Yskes R (2016) Smoking Cessation for People Living With HIV/AIDS: A Literature Review and Synthesis. *Nicotine Tob Res* 18: 2177-2184.
24. Marzolini C, Back D, Weber R, Furrer H, Cavassini, et al. (2011) Ageing with HIV: Medication use and risk for potential drug-drug interactions. *J Antimicrob Chemother* 66: 2107-2111.
25. Hughes CA, Tseng A, Cooper R (2015) Managing drug interactions in HIV-infected adults with comorbid illness. *CMAJ* 187: 36-43.
26. HIV/HCV Drug Therapy Guide (2016) UHN- Toronto General Hospital, Immunodeficiency Clinic. HIV/HCV Drug Therapy Guide, Canada.
27. Loutfy MR, Sherr L, Sonnenberg-Schwan U, Walmsley SL, Johnson M, et al. (2013) Caring for women living with HIV: Gaps in the evidence. *J Int AIDS Soc* 16: 18509.
28. de Pommerol M, Hessamfar M, Lawson-Ayayi S, Neau D, Geffard S, et al. (2011) Menopause and HIV infection: Age at onset and associated factors, ANRS CO₃ Aquitaine cohort. *Int J STD AIDS* 22: 67-72.
29. Ferreira CE, Pinto-Neto AM, Conde DM, Costa-Paiva L, Morais SS, et al. (2007) Menopause symptoms in women infected with HIV: Prevalence and associated factors. *Gynecol Endocrinol* 23: 198-205.
30. Yin MT, Zhang CA, McMahon DJ, Ferris DC, Irani D, et al. (2012) Higher rates of bone loss in postmenopausal HIV-infected women: A longitudinal study. *J Clin Endocrinol Metab* 97: 554-562.
31. Looby SED (2012) Menopause-Associated Metabolic Manifestations and Symptomatology in HIV Infection: A Brief Review with Research Implications. *J Assoc Nurses AIDS Care* 23: 195-203.
32. Linsk NL (2000) HIV among older adults: Age-specific issues in prevention and treatment. *AIDS Read* 10: 430-440.
33. Meditz AL, Moreau KL, MaWhinney S, Gozansky WS, Melander K, et al. (2012) CCR₅ expression is elevated on endocervical CD4+ T cells in healthy postmenopausal women. *J Acquir Immune Defic Syndr* 59: 221-228.
34. Greene M, Covinsky KE, Valcour V, Miao Y, Madamba J, et al. (2015) Geriatric Syndromes in Older HIV-Infected Adults. *J Acquir Immune Defic Syndr* 69: 161-167.
35. Foebel AD, Hirdes JP, Boodram C, Lemick R, Tai JW, et al. (2016) Comparing the care needs of people living with and without HIV in Canadian home and long-term care settings. *Can Commun Dis Rep* 42: 52-56.
36. Siegler EL, Brennan-Ing M (2017) Adapting Systems of Care for People Aging With HIV. *J Assoc Nurses AIDS Care* 28: 698-707.
37. Emler CA (2017) Stigma in an Aging Context. *Interdiscip Top Gerontol Geriatr* 42: 144-158.
38. Emler CA, Brennan DJ, Brennenstuhl S, Rueda S, Hart TA, et al. (2013) Protective and risk factors associated with stigma in a population of older adults living with HIV in Ontario, Canada. *AIDS Care* 25: 1330-1339.
39. Casey House (2020) Compassionate Care in a Changing Landscape: HIV/AIDS and Long-Term Care. Casey House, Toronto, Canada.



- Advances In Industrial Biotechnology | ISSN: 2639-5665
- Advances In Microbiology Research | ISSN: 2689-694X
- Archives Of Surgery And Surgical Education | ISSN: 2689-3126
- Archives Of Urology
- Archives Of Zoological Studies | ISSN: 2640-7779
- Current Trends Medical And Biological Engineering
- International Journal Of Case Reports And Therapeutic Studies | ISSN: 2689-310X
- Journal Of Addiction & Addictive Disorders | ISSN: 2578-7276
- Journal Of Agronomy & Agricultural Science | ISSN: 2689-8292
- Journal Of AIDS Clinical Research & STDs | ISSN: 2572-7370
- Journal Of Alcoholism Drug Abuse & Substance Dependence | ISSN: 2572-9594
- Journal Of Allergy Disorders & Therapy | ISSN: 2470-749X
- Journal Of Alternative Complementary & Integrative Medicine | ISSN: 2470-7562
- Journal Of Alzheimers & Neurodegenerative Diseases | ISSN: 2572-9608
- Journal Of Anesthesia & Clinical Care | ISSN: 2378-8879
- Journal Of Angiology & Vascular Surgery | ISSN: 2572-7397
- Journal Of Animal Research & Veterinary Science | ISSN: 2639-3751
- Journal Of Aquaculture & Fisheries | ISSN: 2576-5523
- Journal Of Atmospheric & Earth Sciences | ISSN: 2689-8780
- Journal Of Biotech Research & Biochemistry
- Journal Of Brain & Neuroscience Research
- Journal Of Cancer Biology & Treatment | ISSN: 2470-7546
- Journal Of Cardiology Study & Research | ISSN: 2640-768X
- Journal Of Cell Biology & Cell Metabolism | ISSN: 2381-1943
- Journal Of Clinical Dermatology & Therapy | ISSN: 2378-8771
- Journal Of Clinical Immunology & Immunotherapy | ISSN: 2378-8844
- Journal Of Clinical Studies & Medical Case Reports | ISSN: 2378-8801
- Journal Of Community Medicine & Public Health Care | ISSN: 2381-1978
- Journal Of Cytology & Tissue Biology | ISSN: 2378-9107
- Journal Of Dairy Research & Technology | ISSN: 2688-9315
- Journal Of Dentistry Oral Health & Cosmesis | ISSN: 2473-6783
- Journal Of Diabetes & Metabolic Disorders | ISSN: 2381-201X
- Journal Of Emergency Medicine Trauma & Surgical Care | ISSN: 2378-8798
- Journal Of Environmental Science Current Research | ISSN: 2643-5020
- Journal Of Food Science & Nutrition | ISSN: 2470-1076
- Journal Of Forensic Legal & Investigative Sciences | ISSN: 2473-733X
- Journal Of Gastroenterology & Hepatology Research | ISSN: 2574-2566
- Journal Of Genetics & Genomic Sciences | ISSN: 2574-2485
- Journal Of Gerontology & Geriatric Medicine | ISSN: 2381-8662
- Journal Of Hematology Blood Transfusion & Disorders | ISSN: 2572-2999
- Journal Of Hospice & Palliative Medical Care
- Journal Of Human Endocrinology | ISSN: 2572-9640
- Journal Of Infectious & Non Infectious Diseases | ISSN: 2381-8654
- Journal Of Internal Medicine & Primary Healthcare | ISSN: 2574-2493
- Journal Of Light & Laser Current Trends
- Journal Of Medicine Study & Research | ISSN: 2639-5657
- Journal Of Modern Chemical Sciences
- Journal Of Nanotechnology Nanomedicine & Nanobiotechnology | ISSN: 2381-2044
- Journal Of Neonatology & Clinical Pediatrics | ISSN: 2378-878X
- Journal Of Nephrology & Renal Therapy | ISSN: 2473-7313
- Journal Of Non Invasive Vascular Investigation | ISSN: 2572-7400
- Journal Of Nuclear Medicine Radiology & Radiation Therapy | ISSN: 2572-7419
- Journal Of Obesity & Weight Loss | ISSN: 2473-7372
- Journal Of Ophthalmology & Clinical Research | ISSN: 2378-8887
- Journal Of Orthopedic Research & Physiotherapy | ISSN: 2381-2052
- Journal Of Otolaryngology Head & Neck Surgery | ISSN: 2573-010X
- Journal Of Pathology Clinical & Medical Research
- Journal Of Pharmacology Pharmaceutics & Pharmacovigilance | ISSN: 2639-5649
- Journal Of Physical Medicine Rehabilitation & Disabilities | ISSN: 2381-8670
- Journal Of Plant Science Current Research | ISSN: 2639-3743
- Journal Of Practical & Professional Nursing | ISSN: 2639-5681
- Journal Of Protein Research & Bioinformatics
- Journal Of Psychiatry Depression & Anxiety | ISSN: 2573-0150
- Journal Of Pulmonary Medicine & Respiratory Research | ISSN: 2573-0177
- Journal Of Reproductive Medicine Gynaecology & Obstetrics | ISSN: 2574-2574
- Journal Of Stem Cells Research Development & Therapy | ISSN: 2381-2060
- Journal Of Surgery Current Trends & Innovations | ISSN: 2578-7284
- Journal Of Toxicology Current Research | ISSN: 2639-3735
- Journal Of Translational Science And Research
- Journal Of Vaccines Research & Vaccination | ISSN: 2573-0193
- Journal Of Virology & Antivirals
- Sports Medicine And Injury Care Journal | ISSN: 2689-8829
- Trends In Anatomy & Physiology | ISSN: 2640-7752

Submit Your Manuscript: <https://www.heraldopenaccess.us/submit-manuscript>